

!!!EMERGING DRUG WARNING!!!


Nitazenes & Medetomidine In NYC's Drug Supply!



Nitazenes

Nitazenes are becoming increasingly common in the drug supply, so read below to learn more!

Nitazenes are opioids up to 10x more potent than fentanyl.



Use Narcan & Call 911 If You Spot These Overdose Signs!

Slow or Shallow Breathing | Loss of Consciousness
Bluish or Grayish Lips & Complexion



Medical Attention May Be Necessary If You Spot These Withdrawal Signs!

Excessive Sweating | Restless Legs | Fever | Dizziness | Flu-Like Symptoms
Blackouts | Anxiety | Panic Attacks


Medetomidine

Medetomidine is becoming increasingly common in the drug supply, so read below to learn more!

Medetomidine is a tranquilizer up to 100x more potent than xylazine.

Medical Attention May Be Necessary If You Spot These Withdrawal Signs!

Fast Heart Rate | Very High Blood Pressure | Nausea & Vomiting | Tremors
Excessive Sweating | Changing Levels of Alertness



Note On Dehydration & Increased Risk of Overdose:

Using medetomidine makes you urinate more, which increases the risk of dehydration, in addition to overdose. **If using, hydrate often!**

IF USING STREET DRUGS...



**USE NARCAN & CALL
911 IF YOU OVERDOSE!**



DON'T USE ALONE!



USE SLOWLY!



STAY WARM!



HYDRATE OFTEN!

USE TEST STRIPS AND DRUG CHECKING LABS!

Check drug supplies for nitazenes and medetomidine using nitazene and medetomidine drug testing strips, OR utilize a drug checking lab with OnPoint NYC, VOCAL-NY, Harlem United, BOOM!Health, or Housing Works (Cylar House).